

MATRICIAL FORMULAE FOR PARTITIONS

F. AICARDI

ABSTRACT. The exponential of the triangular matrix whose entries in the diagonal at distance n from the principal diagonal are all equal to the sum of the inverse of the divisors of n is the triangular matrix whose entries in the diagonal at distance n from the principal diagonal are all equal to the number of partitions of n . A similar result is true for any pair of sequences satisfying a special recurrence.

Let Δ be the infinite triangular matrix having zeroes on and below the main diagonal, and whose values on each parallel to the main diagonal are all equal to:

$$\Delta_{i,i+n} = \bar{\sigma}(n) \quad (i \geq 0, \quad n > 0),$$

where $\bar{\sigma}(n)$ is the sum of the inverses of the divisors of n , for $n > 0$.

Proposition 1. *The matrix*

$$\mathbf{P} = \exp(\Delta)$$

is triangular, with zero below the diagonal, 1 on the diagonal, and its elements on the parallel to the diagonal at distance n from the diagonal are equal to the number $p(n)$ of partitions of the integer n (being $p(0) = 1$):

$$\mathbf{P}_{i,i+n} = p(n) \quad (i \geq 0, \quad n \geq 0).$$

$$\Delta = \begin{pmatrix} 0 & \frac{1}{1} & \frac{3}{2} & \frac{4}{3} & \frac{7}{4} & \frac{6}{5} & \frac{12}{6} & \dots \\ 0 & 0 & \frac{1}{1} & \frac{3}{2} & \frac{4}{3} & \frac{7}{4} & \frac{6}{5} & \dots \\ 0 & 0 & 0 & \frac{1}{1} & \frac{3}{2} & \frac{4}{3} & \frac{7}{4} & \dots \\ 0 & 0 & 0 & 0 & \frac{1}{1} & \frac{3}{2} & \frac{4}{3} & \dots \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{1} & \frac{3}{2} & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{1} & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad \mathbf{P} = \begin{pmatrix} 1 & 1 & 2 & 3 & 5 & 7 & 11 & \dots \\ 0 & 1 & 1 & 2 & 3 & 5 & 5 & \dots \\ 0 & 0 & 1 & 1 & 2 & 3 & 5 & \dots \\ 0 & 0 & 0 & 1 & 1 & 2 & 3 & \dots \\ 0 & 0 & 0 & 0 & 1 & 1 & 2 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

We prove in the next section a more general result. As byproduct, we find a formula (see eq.(3)) for the number of partitions of n as sum of the products of the function $\bar{\sigma}$ over all elements of each partition, and a formula for $\bar{\sigma}$ as combination of the products of the number of partitions over all elements of each partition (see eq.(6)).

Generalisation of Proposition 1.

Let $\{s_n\}_{n=0}^{\infty}$ be any infinite sequence, satisfying $s_0 = 0$.

Let t_n be the infinite sequence obtained by the following recurrence:

$$(1) \quad t_0 = 1, \quad t_n = \frac{1}{n} \sum_{k=0}^{n-1} s_{n-k} t_k$$

Consider the infinite triangular matrix $\mathbf{S}$ having zeroes on the diagonal and below the diagonal, and whose value on each parallel to the diagonal and at distance n from the diagonal are all equal to:

$$\mathbf{S}_{i,i+n} = s_n/n \quad (i \geq 0, \quad n > 0).$$

Theorem 1. *The matrix*

$$\mathbf{T} = \exp(\mathbf{S})$$

is triangular, with zero below the diagonal, 1 on the diagonal, and the values of its elements on each parallel to the diagonal at distance n from the diagonal are equal to t_n :

$$\mathbf{T}_{i,i+n} = t_n \quad (i \geq 0, \quad n \geq 0).$$

Example 1. Proposition 1 is a particular case of Theorem 1. Indeed, $\bar{\sigma}(k) = \sigma(k)/k$, $\sigma(k)$ being the sum of all divisors of k . The recurrence

$$np(n) = \sum_{k=0}^{n-1} \sigma(n-k)p(k) \equiv \sum_{h=1}^n \sigma(h)p(n-h)$$

is proved, for instance in [1]. (A proof can be done in a similar way as we do for the recurrence of the next example).

Example 2. Let $p^o(n)$ be the number of partitions of n into **odd** integers,

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	...
$p^o(n)$	1	1	2	2	3	4	5	6	8	10	12	15	18	22	27	32	38	46	54	64	...

and σ_k^o be the sum of **odd** divisors of k :

k	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	...
$\sigma^o(k)$	1	1	4	1	6	4	8	1	13	6	12	4	14	8	24	1	18	13	20	6	...

Lemma.

$$np^o(n) = \sum_{k=0}^{n-1} \sigma^o(n-k)p^o(k) \equiv \sum_{h=1}^n \sigma^o(h)p^o(n-h).$$

Proof of Lemma. Consider all partitions of n into odd integers

$$n = o_1 + o_2 + \cdots + o_r, \quad o_i \text{ odd},$$

and sum all these equalities: we obtain at left $np^o(n)$, and at right

$$\sum_{i=1}^n o_i \mu_i, \quad \mu_i \geq 0,$$

μ_i being the number of times the odd o_i appears in total inside all partitions of n into odd integers. We have

$$\mu_i = \sum_{m=1}^{n/o_i} p^o(n - mo_i).$$

Indeed, there are possibly several copies of o_i in a partition: $p^o(n - o_i)$ is the number of those containing at least one copy of o_i , and then we count in this summand the first copy of the o_i in all partitions; $p^o(n - 2o_i)$ is the number of partitions containing at least 2 copies of o_i , and then we count in this summand the second copy of o_i in all partitions, and so on. We obtain:

$$np^o(n) = \sum_{i=1}^n o_i \sum_{m=1}^{n/o_i} p^o(n - mo_i).$$

Writing $mo_i = h$, we get

$$np^o(n) = \sum_{h=1}^n \sum_{o_i|h} o_i p^o(n - h),$$

the second sum being extended to all odd integers dividing h . Then

$$p^o(n) = \frac{1}{n} \sum_{h=1}^n \sigma^o(h) p^o(n - h) \equiv \frac{1}{n} \sum_{k=0}^{n-1} \sigma^o(n - k) p^o(k).$$

□

Define the matrix Σ^o whose elements are zero on and below the diagonal and

$$\Sigma_{i,i+n}^o = \sigma^o(n)/n \quad (i \geq 0, \quad n > 0).$$

By Theorem 1, the matrix $\mathbf{P}^o = \exp(\Sigma^o)$ is triangular and satisfies:

$$\mathbf{P}_{i,i+n}^o = p^o(n), \quad (i \geq 0, \quad n \geq 0).$$

Example 3. Let $p^q(n)$ be the number of partitions of n into integers non divisible by the prime q , and σ_k^q be the sum of divisors of k non divisible by q :

Like in the above lemma, we prove

$$np^q(n) = \sum_{h=1}^n \sigma^q(h) p^q(n - h).$$

Define the matrix Σ^q whose elements are zero on and below the diagonal and

$$\Sigma_{i,i+n}^q = \sigma^q(n)/n \quad (i \geq 0, \quad n > 0).$$

By Theorem 1, the matrix $\mathbf{P}^q = \exp(\Sigma^q)$ is triangular and satisfies:

$$\mathbf{P}_{i,i+n}^q = p^q(n) \quad (i \geq 0, \quad n \geq 0).$$

Proof of Theorem 1

The exponential of $\mathbf{S}$ is given by:

$$(2) \quad \exp(\mathbf{S}) = \mathbf{E} + \mathbf{S} + \frac{1}{2!}\mathbf{S}^2 + \frac{1}{3!}\mathbf{S}^3 + \frac{1}{4!}\mathbf{S}^4 + \dots$$

where $\mathbf{E}$ is the identity matrix.

Consider the sequence τ_n , defined as the first row of the matrix $\mathbf{T} = \exp(\mathbf{S})$. We will prove that $\mathbf{T}_{i,i+n} = \tau_n$, and τ_n satisfies the recurrence (1).

We use in the following the notation $s(n)$ instead of s_n in order to avoid many subscript indices. We thus have:

$$\mathbf{S}_{i,k} = s(k-i)/(k-i), \quad \mathbf{S}_{0,k} = s(k)/k.$$

We calculate the powers of $\mathbf{S}$.

The square $\mathbf{S}^2$ is equal to:

$$\mathbf{S}_{i,j}^2 = \sum_{k=0}^n \mathbf{S}_{i,k} \mathbf{S}_{k,j} = \sum_{i < k < j} \frac{s(k-i)s(j-k)}{(k-i)(j-k)},$$

being $\mathbf{S}_{i,k} = 0$ for $k \leq i$. Hence $\mathbf{S}_{i,j}^2 = 0$ for $j < i + 2$.

The matrix $\mathbf{S}^3$ is equal to:

$$\mathbf{S}_{i,j}^3 = \sum_{k=0}^n \mathbf{S}_{i,k} \mathbf{S}_{k,j}^2 = \sum_{k=0}^n \mathbf{S}_{i,k} \sum_{h=0}^n \mathbf{S}_{k,h} \mathbf{S}_{h,j} = \sum_{i < k < h < j} \frac{s(k-i)s(h-k)s(j-h)}{(k-i)(h-k)(j-h)},$$

being $\mathbf{S}_{i,j}^2 = 0$ for $j \leq i + 1$. Hence $\mathbf{S}_{i,j}^3 = 0$ for $j < i + 3$.

We obtain, for the r -th power of $\mathbf{S}$:

$$\mathbf{S}_{i,j}^r = \sum_{i=k_0 < k_1 < k_2 < \dots < k_r=j} \prod_{m=1}^r \frac{s(k_m - k_{m-1})}{(k_m - k_{m-1})},$$

which is zero if $j < i + r$.

The first elements of the sequence $\tau_0, \dots, \tau_n$ are the first elements of the first row of the matrix $\sum_{r=0}^n \frac{1}{r!} (\mathbf{S})^r$, since the elements at distance lower than m over the diagonal of $\mathbf{S}^m$ are zero, as we have seen.

The element $\mathbf{T}_{0,n}$ of the first row of the matrix $\mathbf{E} + \sum_{r=1}^n \frac{1}{r!} \mathbf{S}^r$ are therefore, for $n > 0$

$$\tau_n = \left\{ \mathbf{E} + \sum_{r=1}^n \frac{1}{r!} \mathbf{S}^r \right\}_{0,n} = \sum_{r=1}^n \frac{1}{r!} \sum_{0 < k_1 < k_2 < \dots < k_r=n} \prod_{m=1}^r \frac{s(k_m - k_{m-1})}{(k_m - k_{m-1})},$$

and $\tau_0 = 1$. For every $r = 1 \dots n$, the r positive integers k_m satisfy

$$\sum_{m=1}^r (k_m - k_{m-1}) = n;$$

hence the sum is extended to all partitions of n containing exactly r elements $n = h_1, \dots, h_r$. However, for every partition, the same product $\prod_{m=1}^r s(h_m)/(h_m)$ corresponds to a number of cases with different k_m depending on the ordering of the h_m . This number of different cases is equal to

$$r! / \prod_{m=1}^r \rho_m!,$$

where ρ_m is the multiplicity of h_m in the partition.

Hence we obtain

Proposition 2. *The elements τ_n , for $n > 0$, are equal to:*

$$(3) \quad \tau_n = \sum_{r=1}^n \sum_{X_r(n)} \prod_{m=1}^r \frac{s(h_m)}{h_m \rho_m!}$$

where $X_r(n)$ is the set of partitions of n into exactly r summands h_m , and ρ_m is the multiplicity of h_m .

The formula expressing the elements of $\mathbf{S}^r$ shows that $\mathbf{S}_{i,j}^r$ depends only on the difference $(j - i)$, and therefore the elements $\mathbf{T}_{i,j}$ are equal on each parallel at distance r from the diagonal, where $j = i + r$.

The proof of Theorem 2 is now completed proving that formula (3) satisfies the recurrence:

$$(4) \quad \tau_N = \frac{1}{N} \sum_{n=0}^{N-1} s(N - n) \tau_n.$$

We write the left member of (4) by substituting the terms τ_n given by formula (3):

$$(5) \quad \frac{1}{N} \left(s(N) + \sum_{n=1}^{N-1} s(N - n) \sum_{r=1}^n \sum_{X_r(n)} \prod_{m=1}^r \frac{s(h_m)}{h_m \rho_m!} \right).$$

Each summand in the expression from any τ_n , $0 < n < N$, is the product of r factors $s(h_m)$, such that $\sum_{m=1}^r h_m \rho(m) = n$. In expression (5) it is multiplied by $s(N - n)$ and becomes therefore the product of $r + 1$ factors $s(h_m)$ such that their sum is N , as in the expression of τ_N by (3).

Consider now any one of such terms, containing $\prod_{m=1}^{r+1} s(h_m)$, in which one factor, say $s(h_{m^*})$, is $s(N - n)$.

If h_{m^*} is different from the other r h_m , the contribution of the corresponding term is equal to

$$\frac{1}{N} \frac{1}{\prod_{m=1}^r h_m \rho_m!} = \frac{1}{N} \frac{1}{\prod_{m=1}^{r+1} h_m \rho_m! / h_{m^*}}.$$

If h_{m^*} is equal to one of the other h_m , i.e., in the product has multiplicity $\rho_m > 1$, the coefficient of the corresponding term is equal to

$$\frac{1}{N} \frac{1}{\prod_{m=1}^{r+1} h_m \rho_m! / (h_{m^*} \rho_{m^*})},$$

since $s(h_{m^*=N-n})$ had multiplicity $\rho_{m^*} - 1$ in the original term in τ_n .

If among the $r + 1$ integers h_m , $d \leq r$ integers h_m^* are different, in correspondence of the same product we are considering, there will be in τ_N the contribution of d terms, namely for every value of n such that $N - n = h_{m^*}$.

The coefficient of the terms $\prod_{m=1}^{r+1} s(h_m)$ is thus

$$\frac{1}{N} \sum_{k=1}^d \frac{1}{\prod_{m=1}^{r+1} h_m \rho_m! / (h_{m_k^*} \rho_{m_k^*})},$$

5

which is equal to

$$\frac{\sum_{k=1}^d h_{m_k^*} \rho_{m_k^*}}{N \prod_{m=2}^{r+1} h_m \rho_m!} = \frac{N}{N \prod_{m=1}^{r+1} h_m \rho_m!} = \frac{1}{\prod_{m=1}^{r+1} h_m \rho_m!}.$$

Hence we obtain from eq. (5), denoting $q = r + 1$:

$$\tau_N = \frac{s(N)}{N} + \sum_{r=1}^{N-1} \sum_{X_{r+1}(N)} \prod_{m=1}^{r+1} \frac{s(h_m)}{h_m \rho_m!} = \frac{s(N)}{N} + \sum_{q=2}^N \sum_{X_q(N)} \prod_{m=1}^q \frac{s(h_m)}{h_m \rho_m!},$$

i.e., the expression of τ_N by eq. (3). □

Inverse formula

We see now how the sequence s_n can be written in terms of the sequence t_n satisfying the recurrence (2).

Proposition 3. *The following identity holds:*

$$(6) \quad \frac{s_n}{n} = \sum_{r=1}^n (-1)^{r-1} (r-1)! \sum_{X_r(n)} \prod_{m=1}^r \frac{t_{h_m}}{\rho_m!}$$

Proof.

From the identity $\mathbf{T} = \exp(\mathbf{S})$ we obtain

$$\mathbf{S} = \ln \mathbf{T} = \ln(\mathbf{E} + (\mathbf{T} - \mathbf{E})) = \sum_{r \geq 1} (-1)^{r-1} \frac{1}{r} (\mathbf{T} - \mathbf{E})^r.$$

The first row of matrix $\mathbf{T} - \mathbf{E}$ contains, but the initial elements which is zero, the elements of the sequence t_n . Hence we calculate the elements of $(\mathbf{T} - \mathbf{E})^r$, as in the proof of Theorem 1, and we add them with coefficients $(-1)^{r+1}/r$, to obtain the sequence s_n . □

REFERENCES

- [1] E. Grosswald, *Topics from the theory of numbers*, Macmillan, New York, 1966, p. 226.

(F. Aicardi) SISTIANA 56 PR (TRIESTE), ITALY

E-mail address: aicardi@sissa.it